

Distributed Algorithms 2021


10 Sinkless orientation

This week's plan

- **Topic:** complexity of *sinkless orientation*
 - task: high-degree nodes must have outdegree ≥ 1
 - possible in $O(\log n)$ rounds, not in $o(\log n)$ rounds
- **Video:** *why* do we care about this?
 - e.g. hardness of graph coloring
- **Today:** how to *prove* it?
 - round elimination & fixed points

Sinkless orientation

- **Labels:** $\{ 0, 1 \}$
 - 0 = *"edge oriented away from the active node"*
 - 1 = *"edge oriented towards the active node"*
- **Active:** $[0, ?, ?]$
 - *"at least one outgoing edge"*
- **Passive:** $[1, ?, ?]$
 - *"at least one outgoing edge"*



active:
[0, ?, ?]
passive:
[1, ?, ?]

Sinkless orientation: O, I

- active: $[O, ?, ?]$
- passive: $[I, ?, ?]$

Output problem: $\{O\}, \{I\}, \{O,I\}$

- active: $[\{I\}, ?, ?]$
- passive: $[\{O\}, ?, ?]$ or $[\{O,I\}, ?, ?]$

Maximal problem: $\{I\}, \{O,I\}$

- active: $[\{I\}, \{O,I\}, \{O,I\}]$
- passive: $[\{O,I\}, ?, ?]$

Sinkless orientation: O, I

- active: [$O, ?, ?$]
- passive: [$I, ?, ?$]

Output problem: $\{O\}, \{I\}, \{O,I\}$


- active: [$\{I\}, ?, ?$]
- passive: [$\{O\}, ?, ?$] or [$\{O,I\}, ?, ?$]

Maximal problem: A, B

- active: [A, B, B]
- passive: [$B, ?, ?$]

Output problem

- **Labels:** { A, B }
 - A = *"edge oriented away from the active node"*
 - B = *"edge oriented towards the active node"*
- **Active:** [A, B, B]
 - *"**exactly** one outgoing edge"*
- **Passive:** [B, ?, ?]
 - *"at least one outgoing edge"*



active:
[A, B, B]
passive:
[B, ?, ?]

Starting point: A, B

- active: [A, B, B]
- passive: [B, ?, ?]

Output problem: {A}, {B}, {A,B}

- active: [{B}, ?, ?]
- passive: ...

Maximal problem: {B}, {A,B}

- active: [{B}, {A,B}, {A,B}]
- passive: [{A,B}, ?, ?]

Starting point: A, B

- active: [A, B, B]
- passive: [B, ?, ?]

Output problem: {A}, {B}, {A,B}


- active: [{B}, ?, ?]
- passive: ...

Maximal problem: A, B

- active: [A, B, B]
- passive: [B, ?, ?]

Fixed points

- $X = \mathbf{re}(X)$, and X is not 0-round solvable
- “ X can be solved 1 round faster than X ”
 - contradiction
- One of our assumptions fails — which one?



Fixed points

- $X = \mathbf{re}(X)$, and X is not 0-round solvable
- *X cannot be solved in $o(\log n)$ rounds* in the deterministic PN model
- We can also derive hardness results for deterministic and randomized LOCAL model

Often used like this

- We are interested in problem X
- Find a suitable *relaxation* Y of X
 - problem Y is at most as hard as X
 - problem Y is nontrivial
- Show that $Y = \text{re}(Y)$ or $Y = \text{re}(\text{re}(Y))$
 - Y cannot be solved fast
 - X cannot be solved fast

Sinkless and sourceless

- **Labels:** $\{ 0, 1 \}$
 - 0 = *"edge oriented away from the active node"*
 - 1 = *"edge oriented towards the active node"*
- **Active:** $[0, 1, ?]$
 - *"at least one outgoing and one incoming edge"*
- **Passive:** $[1, 0, ?]$
 - *"at least one outgoing and one incoming edge"*